

POSITIVE DEHN TWIST EXPRESSIONS FOR SOME ELEMENTS OF FINITE ORDER IN THE MAPPING CLASS GROUP

YUSUF Z. GURTAS

ABSTRACT. Positive Dehn twist products for some elements of finite order in the mapping class group of a 2-dimensional closed, compact, oriented surface Σ_g , which are rotations of Σ_g through $2\pi/p$, are presented. The homeomorphism invariants of the resulting simply connected symplectic 4-manifolds are computed.

INTRODUCTION

Positive Dehn twist expressions for a new set of involutions in the mapping class group of 2-dimensional closed, compact, oriented surfaces were presented in [1]. In a later article the idea was extended to bounded surfaces and from that, the positive Dehn twist expressions for involutions on closed surfaces that are obtained by joining several copies of bounded surfaces together were also obtained, [2]. It is the purpose of this article to present positive Dehn twist expressions for some elements of finite order in M_g , which are rotations through $2\pi/p$ on the surface Σ_g . This is shown in section 2 using the expressions for the involutions described in [1]. In section 3 the homeomorphism invariants of the resulting simply connected symplectic 4-manifolds that are realized as Lefschetz fibrations are computed.

1. REVIEW

Let i represent the hyperelliptic (horizontal) involution and s represent the vertical involution as shown in Figure 1.




FIGURE 1. The vertical and horizontal involutions

1991 *Mathematics Subject Classification*. Primary 57M07; Secondary 57R17, 20F38.

Key words and phrases. low dimensional topology, symplectic topology, mapping class group, Lefschetz fibration .

If i is the horizontal involution on a surface Σ_h and s is the vertical involution on a surface Σ_k , k -even, then let θ be the horizontal involution on the surface Σ_g , where $g = h + k$, obtained as in Figure 2.




FIGURE 2. The involution θ on the surface Σ_{h+k}

Figure 3 shows the cycles that are used in expressing θ as a product of positive Dehn twists which is stated in the next theorem.




FIGURE 3. The cycles used in the expression of θ

Theorem 1.0.1. *The positive Dehn twist expression for the involution θ on the surface Σ_{h+k} shown in Figure 3 is given by*

$$\theta = c_{2i+2} \cdots c_{2h} c_{2h+1} c_{2i} \cdots c_{2i-1} b_0 c_{2h+1} c_{2h} \cdots c_{2i+2} c_1 c_2 \cdots c_{2i} b_1 b_2 \cdots b_{k-1} b_k c_{2i+1}.$$

See [1] for the proof.

The order of the twists is from right to left, i.e., c_{2i+1} is applied first.

2. MAIN RESULTS

It is very easy to define many of the elements of finite order in the mapping class group qualitatively using geometry. In this section we will consider those which are realized as rotations on the surface Σ_{1+p} through an angle of $2\pi/p$, Figure 4.




FIGURE 4. $2\pi/p$ rotation ϕ_p on Σ_{1+p}

It is a very difficult task in general, however, to find a positive Dehn twist expression for a given finite order element in the mapping class group. In this section we will find explicit positive Dehn twist expressions for rotations of the kind described above. To achieve that, we will make use of the expression for the involution θ that is given in Theorem 1.0.1. p represents a positive odd integer throughout this section.




FIGURE 5. Definition of the involutions θ_1^3 and θ_2^3 on Σ_{1+3}

Let ϕ_p denote the $2\pi/p$ rotation on the surface Σ_{1+p} as shown in Figure 4.

Let θ_1^p be the involution defined as 180° rotation about the axis through the center of the surface Σ_{1+p} and the hole number $(p+1)/2$, Figure 5 (a). Let θ_2^p

be the involution defined as 180° rotation about the axis through the center of the surface Σ_{1+p} and the hole number 1, Figure 5 (b).

It's not difficult to see that application of θ_1^p followed by θ_2^p results in $2\pi/p$ rotation about the center of the surface. Application of θ_1^p on the surface gives the result shown in Figure 7 (b), and θ_2^p takes this figure to the result shown in Figure 8 (b).

Note that Figure 8 (b) and Figure 6 (b) are the same, namely the result of successive applications of the involutions θ_1^p and θ_2^p is the same as the action of ϕ_p on the surface Σ_{1+p} . From this we conclude that

$$\phi_p = \theta_2^p \theta_1^p$$

on the surface Σ_{1+p} .




FIGURE 6. $2\pi/3$ rotation ϕ_3 on Σ_{1+3}

Now the question is how to find a positive Dehn twist product for ϕ_p . The answer is simply juxtaposing the positive Dehn twist expressions for θ_1^p and θ_2^p . Therefore, we first need to get those expressions for θ_1^p and θ_2^p using Theorem 1.0.1.

Figure 9 shows the cycles that realize θ_1^3 on the surface Σ_{1+3} using Theorem 1.0.1. Similarly, Figure 10 shows the cycles that realize θ_2^3 on the surface Σ_{1+3} using the same theorem.

According to Theorem 1.0.1 we have



$$\theta_1^3 = c_4^1 c_3^1 c_2^1 c_1^1 b_0^1 c_1^1 c_2^1 c_3^1 c_4^1 b_1^1 b_2^1 c_5^1$$

and

$$\theta_2^3 = c_4^2 c_3^2 c_2^2 c_1^2 b_0^2 c_1^2 c_2^2 c_3^2 c_4^2 b_1^2 b_2^2 c_5^2.$$

Therefore

$$\phi_3 = \theta_2^3 \theta_1^3 = c_4^2 c_3^2 c_2^2 c_1^2 b_0^2 c_1^2 c_2^2 c_3^2 c_4^2 b_1^2 b_2^2 c_5^2 c_4^1 c_3^1 c_2^1 c_1^1 b_0^1 c_1^1 c_2^1 c_3^1 c_4^1 b_1^1 b_2^1 c_5^1.$$


 FIGURE 7. Involution θ_1^3 applied to Σ_{1+3}

 FIGURE 8. Involution θ_2^3 applied to the result from Figure 7



Since $\phi_3^3 = 1$, we have

$$(c_4^2 c_3^2 c_2^2 c_1^2 b_0^2 c_2^2 c_3^2 c_4^2 b_1^2 b_2^2 c_5^2 c_4^1 c_3^1 c_2^1 b_0^1 c_1^1 c_2^1 c_3^1 b_1^1 b_2^1 c_5^1)^3 = 1$$

in M_4 , the mapping class group of the surface Σ_4 .

Next, we will demonstrate the cycles used in the expression for ϕ_3 on a regular genus 4 surface Σ_4 and generalize it. For this, we need an identification between the two surfaces Σ_4 and Σ_{1+3} . Figure 11 shows that identification.

Figure 12 shows the same identification for general case. The hole with no number on the right is the hole in the center on the left.

FIGURE 9. Involution θ_1^3 on Σ_{1+3} and the cycles that realize itFIGURE 10. Involution θ_2^3 on Σ_{1+3} and the cycles that realize it






 FIGURE 11. Identification between the two surfaces Σ_4 and Σ_{1+3}


FIGURE 12. Identification between the two surfaces for general case

Figure 13 shows the cycles in Figure 9 (b) via the identification established in Figure 11. Similarly, Figure 14 shows the cycles in Figure 10 (b) via the identification established in Figure 11.



 FIGURE 13. The cycles realizing θ_1^3 on Σ_4

FIGURE 14. The cycles realizing θ_2^3 on Σ_4

In general, the cycles realizing ϕ_p on the round surface Σ_{1+p} are carried onto the regular surface Σ_{p+1} using the identification in Figure 12.

3. APPLICATIONS

Let Σ_g be the 2-dimensional, closed, compact, oriented surface of genus $g > 0$ and M_g be its mapping class group, the group of isotopy classes of all orientation-preserving diffeomorphisms $\Sigma_g \rightarrow \Sigma_g$. It is a well-known fact that any word in the mapping class group M_g that contains positive exponents only and is equal to the identity element defines a symplectic Lefschetz fibration $X^4 \rightarrow S^2$. In this section we compute the homeomorphism invariants of the symplectic Lefschetz fibrations that are defined by the words $\phi_p^p = 1$, p - odd, in M_{p+1} .

FIGURE 15. Standard generators of $\pi_1(\Sigma_{p+1})$

First, we will show that the 4- manifold X carrying the symplectic Lefschetz fibration structure $X^4 \rightarrow S^2$ is simply connected. A well known fact in theory of Lefschetz fibrations states that $\pi_1(X)$ is isomorphic to the quotient of $\pi_1(\Sigma_{p+1})$

by the normal subgroup generated by the vanishing cycles, the cycles about which positive Dehn twists in the expression $\phi_p^p = 1$ are performed. The vanishing cycles are seen as elements of $\pi_1(\Sigma_{p+1})$ in this section, not as elements of M_{g+1} .

Let $\{\alpha_1, \beta_1, \dots, \beta_{p+1}, \alpha_{p+1}\}$ be the set of standard $2p+2$ generators of $\pi_1(\Sigma_{p+1})$ as shown in Figure 15. We will show that the subgroup of $\pi_1(\Sigma_{p+1})$ generated by the vanishing cycles includes all the generators of $\pi_1(\Sigma_{p+1})$ by showing that each generator is equal to 1 in the quotient group. We will use the elements in $\pi_1(\Sigma_{p+1})$ that are shown in Figure 16 along with the standard generators to express the vanishing cycles as elements of $\pi_1(\Sigma_{p+1})$. The expressions should be read from right to left.




FIGURE 16. $\gamma_i = \alpha_i \alpha_{i+1}^{-1}$, $1 \leq i \leq p$

If $p = 3$ then

$$\begin{aligned}\theta_1^3 &= c_4^1 c_3^1 c_2^1 c_1^1 b_0^1 c_1^1 c_2^1 c_3^1 c_4^1 b_1^1 b_2^1 c_5^1 \\ \theta_2^3 &= c_4^2 c_3^2 c_2^2 c_1^2 b_0^2 c_1^2 c_2^2 c_3^2 c_4^2 b_1^2 b_2^2 c_5^2\end{aligned}$$

and we have the following expressions in $\pi_1(\Sigma_4)$ for the vanishing cycles shown in Figures 13 and 14:

$$\begin{aligned}c_1^1 &= \alpha_1, c_2^1 = \beta_1, c_3^1 = \gamma_1, c_4^1 = \beta_2, c_5^1 = \beta_3 \alpha_3^{-1} \beta_3^{-1} \gamma_2^{-1} \\ b_0^1 &= \beta_4^{-1} \beta_3^{-1} \beta_2^{-1} \beta_1^{-1} \\ b_1^1 &= \alpha_3 \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \\ b_2^1 &= \beta_3 \alpha_3 \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \\ c_1^2 &= \alpha_3, c_2^2 = \beta_3, c_3^2 = \gamma_2, c_4^2 = \beta_2, c_5^2 = \gamma_1^{-1} \beta_1^{-1} \alpha_1 \beta_1 \\ b_0^2 &= \beta_4^{-1} \beta_3^{-1} \beta_2^{-1} \beta_1^{-1} \\ b_1^2 &= \gamma_1 \beta_2 \beta_3 \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2^{-1} \gamma_1^{-1} \beta_1^{-1} \alpha_1 \\ b_2^2 &= \gamma_1 \beta_2 \beta_3 \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2^{-1} \gamma_1^{-1} \beta_1^{-1} \alpha_1 \beta_1\end{aligned}$$

Let $q = \frac{p+1}{2}$, i.e., $p = 2q - 1$. Then for $p \geq 5$, and hence $q \geq 3$,

$$\begin{aligned}(3.0.1) \quad \theta_2^p &= c_4^2 c_3^2 c_2^2 c_1^2 b_0^2 c_1^2 c_2^2 c_3^2 c_4^2 b_1^2 \cdots b_{p-1}^2 c_5^2 \\ \theta_1^p &= c_4^1 c_3^1 c_2^1 c_1^1 b_0^1 c_1^1 c_2^1 c_3^1 c_4^1 b_1^1 \cdots b_{p-1}^1 c_5^1\end{aligned}$$

and we have the following expressions in $\pi_1(\Sigma_{p+1})$ for the vanishing cycles:

θ_1^p cycles:

$$c_1^1 = \alpha_1, c_2^1 = \beta_1, c_3^1 = \gamma_1, c_4^1 = \beta_2$$

$$c_5^1 = \beta_3 \cdots \beta_q \beta_{q+1} \alpha_{q+1}^{-1} \beta_{q+1}^{-1} \gamma_q^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1}$$

$$b_0^1 = \beta_{p+1}^{-1} \beta_p^{-1} \cdots \beta_2^{-1} \beta_1^{-1}$$

$$b_1^1 = \alpha_3 \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1}$$

$$b_{2m}^1 = \beta_3 \cdots \beta_{m+1} \beta_{m+2} \alpha_{m+2} \gamma_{p-m+1}^{-1} \beta_{p-m+1}^{-1} \gamma_{p-m}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1}, q-1 \geq m \geq 1$$

$$b_{2m-1}^1 = \beta_3 \cdots \beta_m \beta_{m+1} \alpha_{m+2} \beta_{p-m+2}^{-1} \gamma_{p-m+1}^{-1} \beta_{p-m+1}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1}, q-1 \geq m \geq 2$$

θ_2^p cycles:

$$c_1^2 = \alpha_{q+1}, c_2^2 = \beta_{q+1}, c_4^2 = \beta_2, c_5^2 = \gamma_1^{-1} \beta_1^{-1} \alpha_1 \beta_1$$

$$c_3^2 = \beta_3 \cdots \beta_{q-1} \beta_q \gamma_q^{-1} \beta_q^{-1} \cdots \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1}$$

$$b_0^2 = \beta_{p+1}^{-1} \beta_p^{-1} \cdots \beta_2^{-1} \beta_1^{-1}$$

$$b_{p-1}^2 = \gamma_1 \beta_2 \beta_3 \cdots \beta_{p-1} \beta_p \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 \beta_1$$

$$b_{p-2}^2 = \gamma_1 \beta_2 \beta_3 \cdots \beta_{p-1} \beta_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1$$

$$b_{2m-1}^2 = \beta_{1+q-m}^{-1} \gamma_{q-m}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2 \beta_3 \cdots \beta_{q+m} \gamma_{q+m+1} \cdots \gamma_{p-1} \gamma_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1, \\ q-2 \geq m \geq 1$$

$$b_{2m}^2 = \gamma_{q-m}^{-1} \beta_{q-m}^{-1} \cdots \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2 \beta_3 \cdots \beta_{q+m+1} \gamma_{q+m+1} \cdots \gamma_{p-1} \gamma_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1, \\ q-2 \geq m \geq 1.$$

All of the vanishing cycles above are equal to 1 in the quotient $\pi_1(\Sigma_{p+1})/N$, where N is the normal subgroup generated by the vanishing cycles. Therefore we can cancel some of the standard generators right away: $\alpha_1, \beta_1, \beta_2, \alpha_{q+1}, \beta_{q+1}$ can be cancelled because they are equal to the cycles $c_1^1, c_2^1, c_4^1, c_1^2, c_2^2$, respectively. Also, α_2 can be cancelled because $c_3^1 = \gamma_1 = \alpha_1 \alpha_2^{-1}$. There remains $2p-4$ elements, which are

$$(3.0.2) \quad \alpha_3, \alpha_4, \dots, \alpha_q, \alpha_{q+2}, \dots, \alpha_p, \alpha_{p+1} \beta_3, \beta_4, \dots, \beta_q, \beta_{q+2} \cdots, \beta_p, \beta_{p+1}.$$

Now,

$$b_1^1 = \alpha_3 \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = 1$$

and

$$b_2^1 = \beta_3 \alpha_3 \gamma_p^{-1} \beta_p^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = 1$$

give

$$\alpha_3 \beta_{p+1}^{-1} = \beta_3 \alpha_3, \text{ i.e., } \beta_3 = \alpha_3 \beta_{p+1}^{-1} \alpha_3^{-1}.$$

Similarly,

$$b_3^1 = \beta_3 \alpha_4 \beta_p^{-1} \gamma_{p-1}^{-1} \beta_{p-1}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = 1$$

and

$$b_4^1 = \beta_3 \beta_4 \alpha_4 \gamma_{p-1}^{-1} \beta_{p-1}^{-1} \gamma_{p-2}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = 1$$

give

$$\beta_3 \beta_4 \alpha_4 = \beta_3 \alpha_4 \beta_p^{-1}, \text{ i.e., } \beta_4 = \alpha_4 \beta_p^{-1} \alpha_4^{-1}.$$

In general

$$b_{2m-1}^1 = \beta_3 \cdots \beta_{m+1} \alpha_{m+2} \beta_{p-m+2}^{-1} \gamma_{p-m+1}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = 1$$

and

$$b_{2m}^1 = \beta_3 \cdots \beta_{m+2} \alpha_{m+2} \gamma_{p-m+1}^{-1} \beta_{p-m+1}^{-1} \gamma_{p-m}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = 1$$

give

$$(3.0.3) \quad \beta_{m+2} = \alpha_{m+2} \beta_{p-m+2}^{-1} \alpha_{m+2}^{-1}, \quad q-1 \geq m \geq 1.$$

A similar conjugation relation can be obtained by looking at the cycles of θ_2^p in pairs as well. The first pair is b_1^2 and b_2^2 :

$$b_1^2 = \beta_q^{-1} \gamma_{q-1}^{-1} \cdots \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2 \cdots \beta_{q+1} \gamma_{q+2} \cdots \gamma_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 = 1$$

and

$$b_2^2 = \gamma_{q-1}^{-1} \beta_{q-1}^{-1} \cdots \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2 \cdots \beta_{q+2} \gamma_{q+2} \cdots \gamma_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 = 1$$

result in

$$\beta_q^{-1}\gamma_{q-1}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+1}=\gamma_{q-1}^{-1}\beta_{q-1}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+1}\beta_{q+2},$$

which can be written as

$$(\gamma_{q-1}^{-1}\beta_{q-1}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+1})^{-1}\beta_q^{-1}(\gamma_{q-1}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+1})=\beta_{q+2}.$$

Similarly, from

$$b_3^2=\beta_{q-1}^{-1}\gamma_{q-2}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+2}\gamma_{q+3}\cdots\gamma_p\beta_{p+1}^{-1}\gamma_p^{-1}\beta_p^{-1}\gamma_{p-1}^{-1}\cdots\gamma_1^{-1}\beta_1^{-1}\alpha_1=1$$

and

$$b_4^2=\gamma_{q-2}^{-1}\beta_{q-2}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+3}\gamma_{q+3}\cdots\gamma_p\beta_{p+1}^{-1}\gamma_p^{-1}\beta_p^{-1}\gamma_{p-1}^{-1}\cdots\gamma_1^{-1}\beta_1^{-1}\alpha_1=1$$

we obtain

$$\beta_{q-1}^{-1}\gamma_{q-2}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+2}=\gamma_{q-2}^{-1}\beta_{q-2}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+2}\beta_{q+3},$$

which can be written as

$$(\gamma_{q-2}^{-1}\beta_{q-2}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+2})^{-1}\beta_{q-1}^{-1}(\gamma_{q-2}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+2})=\beta_{q+3}.$$

In general, from

$$b_{2m-1}^2=\beta_{1+q-m}^{-1}\gamma_{q-m}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+m}\gamma_{q+m+1}\cdots\gamma_p\beta_{p+1}^{-1}\gamma_p^{-1}\beta_p^{-1}\gamma_{p-1}^{-1}\cdots\gamma_1^{-1}\beta_1^{-1}\alpha_1=1$$

and

$$b_{2m}^2=\gamma_{q-m}^{-1}\beta_{q-m}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+m+1}\gamma_{q+m+1}\cdots\gamma_p\beta_{p+1}^{-1}\gamma_p^{-1}\beta_p^{-1}\gamma_{p-1}^{-1}\cdots\gamma_1^{-1}\beta_1^{-1}\alpha_1=1$$

we obtain

$$\beta_{1+q-m}^{-1}\gamma_{q-m}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+m}=\gamma_{q-m}^{-1}\beta_{q-m}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+m}\beta_{q+m+1}$$

which can be written as

$$(3.0.4) \quad \beta_{1+q-m}=(\gamma_{q-m}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+m})\beta_{q+m+1}^{-1}(\gamma_{q-m}^{-1}\beta_{q-m}^{-1}\cdots\gamma_3^{-1}\beta_3^{-1}\gamma_2^{-1}\beta_2\cdots\beta_{q+m})^{-1},$$

$$q - 2 \geq m \geq 1$$

Setting $m = 1$ in 3.0.3 we see that β_3 and β_{p+1}^{-1} are conjugate. On the other hand substituting $m = q - 2$ in 3.0.4 we see that β_3 and β_p^{-1} are conjugate. Likewise setting $m = 2$ in 3.0.3 and $m = q - 3$ in 3.0.4 we see that β_4, β_p^{-1} and β_{p-1}^{-1} are conjugate to each other. Continuing this way we get the following list of triples.

$$\begin{array}{ccc} \beta_3 & \beta_{p+1}^{-1} & \beta_p^{-1} \\ \beta_4 & \beta_p^{-1} & \beta_{p-1}^{-1} \\ \beta_5 & \beta_{p-1}^{-1} & \beta_{p-2}^{-1} \\ \dots & \dots & \dots \\ \beta_q & \beta_{q+3}^{-1} & \beta_{q+2}^{-1} \end{array}$$

Each row in this list contains elements that are conjugate to each other. It is easy now to see that $\beta_3, \beta_4, \dots, \beta_q, \beta_{q+2}^{-1}, \beta_{q+3}^{-1}, \dots, \beta_p^{-1}, \beta_{p+1}^{-1}$ are all conjugate to each other. Finally, setting $m = q - 1$ in 3.0.3 we see that β_{q+1} and β_{q+2}^{-1} are conjugate. Since $c_2^2 = \beta_{q+1} = 1$ we conclude that the second half of the generators in 3.0.2 are all identity in the quotient. Instead of $\beta_{q+1} = 1$ here we could have used $\beta_{p+1} = 1$, which is obtained from the pair of cycles b_{p-1}^2, b_{p-2}^2 :

$$b_{p-1}^2 = \gamma_1 \beta_2 \cdots \beta_{p-1} \beta_p \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 \beta_1 = 1$$

and

$$b_{p-2}^2 = \gamma_1 \beta_2 \cdots \beta_{p-1} \beta_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 = 1$$

give

$$\gamma_1 \beta_2 \cdots \beta_{p-1} \beta_p = \gamma_1 \beta_2 \cdots \beta_{p-1} \beta_p \beta_{p+1}^{-1}, \text{ i.e., } \beta_{p+1} = 1$$

using the fact that $c_2^1 = \beta_1 = 1$.

Therefore, the list of generators reduces to

$$\alpha_3, \alpha_4, \dots, \alpha_q, \alpha_{q+2}, \dots, \alpha_p, \alpha_{p+1}.$$

Before we proceed further let's recall that $\gamma_i = \alpha_i \alpha_{i+1}^{-1}$. Therefore

$$\begin{aligned} \gamma_i \cdots \gamma_j &= \alpha_i \alpha_{i+1}^{-1} \alpha_{i+1} \alpha_{i+2}^{-1} \cdots \alpha_{j-1} \alpha_j^{-1} \alpha_j \alpha_{j+1}^{-1} = \alpha_i \alpha_{j+1}^{-1} \\ \gamma_j^{-1} \cdots \gamma_i^{-1} &= \alpha_{j+1} \alpha_j^{-1} \alpha_j \alpha_{j-1}^{-1} \cdots \alpha_{i+2} \alpha_{i+1}^{-1} \alpha_{i+1} \alpha_i^{-1} = \alpha_{j+1} \alpha_i^{-1} \end{aligned}$$

for $j > i$. Now,

$$b_1^1 = \alpha_3 \beta_{p+1}^{-1} \gamma_p^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = \alpha_3 \gamma_p^{-1} \cdots \gamma_3^{-1} \gamma_2^{-1} = \alpha_3 \alpha_{p+1} \alpha_2^{-1} = \alpha_3 \alpha_{p+1} = 1.$$

In general we have

$$\begin{aligned}
 b_{2m-1}^1 &= \beta_3 \cdots \beta_{m+1} \alpha_{m+2} \beta_{p-m+2}^{-1} \gamma_{p-m+1}^{-1} \cdots \beta_4^{-1} \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} = \alpha_{m+2} \gamma_{p-m+1}^{-1} \cdots \gamma_3^{-1} \gamma_2^{-1} \\
 (3.0.5) \quad &= \alpha_{m+2} \alpha_{p-m+2} \alpha_2^{-1} = \alpha_{m+2} \alpha_{p-m+2} = 1, \quad q-1 \geq m \geq 2.
 \end{aligned}$$

A similar list of products can be obtained using the cycles of θ_2^p as follows:

$$\begin{aligned}
 b_1^2 &= \beta_q^{-1} \gamma_{q-1}^{-1} \cdots \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2 \cdots \beta_{q+1} \gamma_{q+2} \cdots \gamma_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 \\
 &= \gamma_{q-1}^{-1} \cdots \gamma_2^{-1} \gamma_{q+2} \cdots \gamma_p \gamma_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \\
 &= \gamma_{q-1}^{-1} \cdots \gamma_2^{-1} \gamma_{q+1}^{-1} \cdots \gamma_1^{-1} \\
 &= \alpha_q \alpha_2^{-1} \alpha_{q+2} \alpha_1^{-1} \\
 &= \alpha_q \alpha_{q+2} = 1
 \end{aligned}$$

In general

$$\begin{aligned}
 b_{2m-1}^2 &= \beta_{1+q-m}^{-1} \gamma_{q-m}^{-1} \cdots \gamma_3^{-1} \beta_3^{-1} \gamma_2^{-1} \beta_2 \cdots \beta_{q+m} \gamma_{q+m+1} \cdots \gamma_p \beta_{p+1}^{-1} \gamma_p^{-1} \beta_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \beta_1^{-1} \alpha_1 \\
 &= \gamma_{q-m}^{-1} \cdots \gamma_2^{-1} \gamma_{q+m+1} \cdots \gamma_p \gamma_p^{-1} \gamma_{p-1}^{-1} \cdots \gamma_1^{-1} \\
 &= \gamma_{q-m}^{-1} \cdots \gamma_2^{-1} \gamma_{q+m}^{-1} \cdots \gamma_1^{-1} \\
 &= \alpha_{q-m+1} \alpha_2^{-1} \alpha_{q+m+1} \alpha_1^{-1} \\
 (3.0.6) \quad &= \alpha_{q-m+1} \alpha_{q+m+1} = 1
 \end{aligned}$$

$$q-2 \geq m \geq 1$$

The product relations in 3.0.5 and 3.0.6 can be summarized as

$$\begin{aligned}
 \alpha_3 \alpha_{p+1} &= 1 \\
 \alpha_4 \alpha_p &= 1 \\
 &\vdots \\
 \alpha_q \alpha_{q+3} &= 1 \\
 \alpha_{q+1} \alpha_{q+2} &= 1
 \end{aligned}$$

and

$$\begin{aligned}
 \alpha_q \alpha_{q+2} &= 1 \\
 \alpha_{q-1} \alpha_{q+3} &= 1 \\
 &\vdots \\
 \alpha_4 \alpha_{p-1} &= 1 \\
 \alpha_3 \alpha_p &= 1
 \end{aligned}$$

respectively. Using $c_1^2 = \alpha_{q+1} = 1$ we see that $\alpha_{q+2} = 1$. If $\alpha_{q+2} = 1$, then $\alpha_q = 1$ and if $\alpha_q = 1$, then $\alpha_{q+3} = 1$, etc. Continuing this way we can conclude that $\alpha_3 = \alpha_4 = \dots = \alpha_{p+1} = 1$ in the quotient $\pi_1(\Sigma_{p+1})/N$, which is isomorphic to $\pi_1(X)$, and this concludes the proof.

We do not know a topological argument to determine what those simply connected symplectic 4-manifolds are but we can compute their homeomorphism invariants using the symplectic Lefschetz fibration structure that they support. This will be accomplished by computing the Euler characteristics and the signatures of those Lefschetz fibrations.

Another fact that we will borrow from the theory of Lefschetz fibrations is the fact that the Euler characteristic of such a fibration $X^4 \rightarrow S^2$ is given by the formula

$$\chi(X) = 4 - 4g + s,$$

where s is the number of singular fibers, i.e., the number of vanishing cycles.

From 3.0.1 we see that each of θ_1^p and θ_2^p consists of $p + 9$ cycles. Therefore $\phi_p^p = (\theta_2^p \theta_1^p)^p$ consists of

$$2p(p + 9)$$

cycles and the Euler characteristic is found to be

$$\begin{aligned} \chi(X) &= 4 - 4(p + 1) + 2p(p + 9) \\ &= 2p^2 + 14p \\ &= 2p(p + 7). \end{aligned}$$

In order to compute the signature $\sigma(X)$ we wrote a Matlab program using the algorithm described in [3]. The computations that we have done using this program point to the closed formula

$$\sigma(X) = -12p$$

for the signature. Taking this formula for granted we can conclude that

$$\begin{aligned} c_1^2(X) &= 3\sigma(X) + 2\chi(X) \\ &= 3(-12p) + 2 \cdot 2p(p + 7) \\ &= 4p(p - 2) \end{aligned}$$

and

$$\begin{aligned} \chi_h(X) &= \frac{\sigma(X) + \chi(X)}{4} \\ &= \frac{-12p + 2p(p + 7)}{4} \\ &= \frac{1}{2}p(p + 1). \end{aligned}$$

$\chi_h(X)$ in the above computation makes sense because X has almost complex structure.

4. APPENDIX

We include the actual outputs of signature computations for some small values of p .

$p = 3 :$

0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, -1, -1, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, -1, 0, -1, -1, -1, -1, 0, 0, 0, 0, 0

$p = 5 :$

0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, -1, -1, 0, 0, 0, 0, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, -1, 0, 0, 0, -1, -1, -1, -1, 0, 0, 0, 0, 0

$p = 7 :$

0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, -1, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0, 0

$p = 9 :$

0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, -1, -1, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, -1, 0, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, 0, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0,
 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, -1, -1, -1, 0, 0, 0, 0, 0, 0, 0, -1, -1, -1, -1, -1, 0, 0, 0, 0, 0

REFERENCES

1. Y.Z. Gurtas, *Positive Dehn Twist Expressions for some New Involutions in the Mapping Class Group M_n* , preprint, arXiv:math.GT/0404310.
2. Y.Z. Gurtas, *Positive Dehn Twist Expressions for some New Involutions in the Mapping Class Group M_n* , preprint, arXiv:math.GT/0404311.
3. B.Ozbagci, *Signatures of Lefschetz fibrations*, Pacific Journal of Mathematics, Vol. **202** 1 (2002), 99-118.

DEPARTMENT OF MATHEMATICS, SUFFOLK CCC, SELDEN, NY, USA
E-mail address: gurtasy@sunysuffolk.edu